

Human CTGF/CCN2 Protein

Cat. No. CGF-HM201



Description

Source	Recombinant Human CTGF/CCN2 Protein is expressed from HEK293 with hFc tag at the N-Terminus. It contains Gln27-Ala349.
Accession	Q5M8T4
Molecular Weight	The protein has a predicted MW of 61.77 kDa. Due to glycosylation, the protein migrates to 50-55 kDa and 62-72 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Connective tissue growth factor (CTGF) is a member of the CCN matricellular protein family, consisting of four domains, that regulates the signaling of other growth factors and promotes kidney fibrosis. CTGF can simultaneously interact with several factors with its four domains. The microenvironment differs depending on the types of cells and tissues and differentiation stages of these cells.


Assay Data

Bis-Tris PAGE



Human CTGF on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human CTGF is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human CTGF, hFc Tag ELISA
0.1µg Human CTGF, hFc Tag Per Well



Immobilized Human CTGF, hFc Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-CTGF Antibody, hFc Tag with the EC50 of 25.8ng/ml determined by ELISA (QC Test).