
Biotinylated SARS-COV-2 Spike S Trimer Protein

Cat. No. COV-VM4SSB

Description	
Source	Recombinant Biotinylated SARS-COV-2 Spike S Trimer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Val16-Glu1188.
Accession	QHD43416.1
Molecular Weight	The protein has a predicted MW of 136.6 kDa. Due to glycosylation, the protein migrates to 140-170 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction.The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Assay Data

Bis-Tris PAGE




Biotinylated SARS-COV-2 Spike S Trimer on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

KVCJUS

ELISA Data

Biotinylated SARS-COV-2 Spike S (Trimer), His Tag ELISA 0.2µg Biotinylated SARS-COV-2 Spike S (Trimer), His Tag Per Well



Immobilized Biotinylated SARS-COV-2 Spike S (Trimer), His Tag at 2μ g/ml (100μ I/well) on the streptavidin precoated plate (5μ g/ml). Dose response curve for Human ACE2, hFc Tag with the EC50 of 5.9ng/ml determined by ELISA.