

Biotinylated SARS-COV-2 Spike S1 Protein

Cat. No. COV-VM5S1B

Description

Source	Recombinant Biotinylated SARS-COV-2 Spike S1 Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus. It contains Gln14-Arg683.
Accession	QHD43416.1
Molecular Weight	The protein has a predicted MW of 102.6 kDa. Due to glycosylation, the protein migrates to 130-140 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 24 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.


Assay Data

Bis-Tris PAGE



Biotinylated SARS-COV-2 Spike S1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated SARS-COV-2 Spike S1 is greater than 95% as determined by SEC-HPLC.

Biotinylated SARS-COV-2 Spike S1 Protein

Cat. No. COV-VM5S1B




Assay Data

ELISA Data

Biotinylated SARS-COV-2 Spike S1, hFc Tag ELISA

0.2µg Biotinylated SARS-COV-2 Spike S1, hFc Tag Per Well



Immobilized Biotinylated SARS-COV-2 Spike S1, hFc Tag at 2µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human ACE2, His Tag with the EC50 of 0.09µg/ml determined by ELISA.