

Canine CRTAM Protein

Cat. No. CRM-DM101

Description

| | |
|-------------------------|---|
| Source | Recombinant Canine CRTAM Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Phe18-Gly287. |
| Accession | E2QWY2 |
| Molecular Weight | The protein has a predicted MW of 31.19 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Tris-Bis PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Tris-Bis PAGE |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Class-I Restricted T Cell-Associated Molecule (CRTAM) is a protein that is expressed after T cell activation. The interaction of CRTAM with its ligand, nectin-like 2 (Nectin2), is required for the efficient production of IL-17, IL-22, and IFN γ by murine CD4 T cells, and it plays a role in optimal CD8 T and NK cell cytotoxicity. CRTAM promotes the pro-inflammatory cytokine profile; therefore, it may take part in the immunopathology of autoimmune diseases such as diabetes type 1 or colitis.

Assay Data

Tris-Bis PAGE



Canine CRTAM on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.