

Cynomolgus CX3CL1/Fractalkine Protein

Cat. No. CX3-CM101

Description

Source	Recombinant Cynomolgus CX3CL1/Fractalkine Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln25-Gln341.
Accession	EHH60415.1
Molecular Weight	The protein has a predicted MW of 34.65 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Fractalkine/CX3C chemokine ligand 1 (CX3CL1) is a chemokine involved in the anticancer function of lymphocytes-mainly NK cells, T cells and dendritic cells. Its increased levels in tumors improve the prognosis for cancer patients, although it is also associated with a poorer prognosis in some types of cancers, such as pancreatic ductal adenocarcinoma.

Assay Data

Bis-Tris PAGE



Cynomolgus CX3CL1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.