Biotinylated Human DNAM-1/CD226 Protein






Description	
Source	Recombinant Biotinylated Human DNAM-1/CD226 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Glu19Asn247.
Accession	Q15762
Molecular Weight	The protein has a predicted MW of 28.9 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by SEC-HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	DNAY

DNAX accessory molecule-1 (DNAM-1), also known as CD226, is a 65 kDa type I transmembrane glycoprotein in the immunoglobulin superfamily.DNAM-1 mediates cellular adhesion to other cells bearing its ligands, CD112 and CD155, and cross-linking DNAM-1 with antibodies causes cellular activation. Furthermore, DNAM-1 can interact with PVR and PVRL2.

Assay Data

Bis-Tris PAGE




Biotinylated Human DNAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS

Assay Data




The purity of Biotinylated Human DNAM-1 is greater than 95% as determined by SEC-HPLC.

ELISA Data


Biotinylated Human DNAM-1, His Tag ELISA

0.1µg Biotinylated Human DNAM-1, His Tag Per Well



Immobilized Biotinylated Human DNAM-1, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate(5µg/ml). Dose response curve for Human CD155, mFc Tag with the EC50 of 11.0ng/ml determined by ELISA (QC Test).

SPR Data



Human CD155, hFc Tag captured on CM5 Chip via Protein A can bind Biotinylated Human DNAM-1, His-Avi Tag with an affinity constant of 264nM as determined in SPR assay (Biacore T200).