

Biotinylated Human DDR1 Protein

Cat. No. DDR-HM4R1B



Description

Source	Recombinant Biotinylated Human DDR1 Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Asp21-Ala417.
Accession	Q08345-1
Molecular Weight	The protein has a predicted MW of 46.93 kDa. Due to glycosylation, the protein migrates to 60-75 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Discoidin domain receptor1 (DDR1) is a collagen activated receptor tyrosine kinase and an attractive anti-fibrotic target. Its expression is mainly limited to epithelial cells located in several organs including skin, kidney, liver and lung. DDR1's biology is elusive, with unknown downstream activation pathways; however, it may act as a mediator of the stromal-epithelial interaction, potentially controlling the activation state of the resident quiescent fibroblasts.


Assay Data

Bis-Tris PAGE



Biotinylated Human DDR1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human DDR1 is greater than 95% as determined by SEC-HPLC.

Biotinylated Human DDR1 Protein


Cat. No. DDR-HM4R1B



Assay Data

ELISA Data

Biotinylated Human DDR1, His Avi Tag ELISA
0.05µg Biotinylated Human DDR1, His Avi Tag Per Well



Immobilized Biotinylated Human DDR1, His Avi Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-DDR1 Antibody, hFc Tag with the EC50 of 7.0ng/ml determined by ELISA.