# **Human DKK1 Protein**

#### Cat. No. DKK-HM201



Description	
Source	Recombinant Human DKK1 Protein is expressed from HEK293 with hFc tag at the C-terminus.
	It contains Thr32-His266.
Accession	O94907
Molecular Weight	The protein has a predicted MW of 52.54 kDa. Due to glycosylation, the protein migrates to 65-80 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

## Formulation and Storage

Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after

optimal storage. Please minimize freeze-thaw cycles.

# Background


Storage

Dickkopf related protein 1 (Dkk-1) is the founding member of the Dickkopf family of proteins that includes Dkk-1, -2, -3, -4, and a related protein, Soggy. Dkk proteins are secreted proteins that contain two conserved cysteinerich domains separated by a linker region. Dkk antagonizes canonical Wnt signaling by inhibiting LRP5/6 interaction with Wnt and by forming a ternary complex with the transmembrane protein KREMEN that promotes internalization of LRP5/6. DKKs play an important role in vertebrate development, where they locally inhibit Wnt regulated processes such as antero-posterior axial patterning, limb development, somitogenesis and eye formation.

reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for

# **Assay Data**

#### Tris-Bis PAGE




Human DKK1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

**SEC-HPLC** 

# KAGTUS

# **Assay Data**



The purity of Human DKK1 is greater than 95% as determined by SEC-HPLC.




### **Assay Data**

**ELISA Data** 

# Human DKK1, hFc Tag ELISA

0.5μg Human DKK1, hFc Tag Per Well



Immobilized Human DKK1, hFc Tag at  $5\mu g/ml$  (100 $\mu l/well$ ) on the plate. Dose response curve for Human LRP-5, mFc Tag with the EC50 of 1.69 $\mu g/ml$  determined by ELISA.