

Cat. No. DLL-HM103B-UL

Description	
Source	Recombinant Biotinylated Human DLL3 Protein (Primary Amine Labeling) is expressed from HEK293 with His tag at the N-Terminus It contains Ala27-Arg490.
Accession	Q9NYJ7-1
Molecular Weight	The protein has a predicted MW of 50.4 kDa. Due to glycosylation, the protein migrates to 52-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC


Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS, 200 mM L-Arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Delta-like protein 3 (DLL3) is a transmembrane protein that belongs to the Delta/Serrate/Lag-2 (DSL) family of Notch ligands. DLL3 inhibits primary neurogenesis. May be required to divert neurons along a specific differentiation pathway. Plays a role in the formation of somite boundaries during segmentation of the paraxial mesoderm (By similarity).

Assay Data


Bis-Tris PAGE



Biotinylated Human DLL3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Biotinylated Human DLL3 is greater than 90% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human DLL3, His Tag ELISA

0.05µg Biotinylated Human DLL3, His Tag Per Well



Immobilized Biotinylated Human DLL3, His Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-DLL3 Antibody, hFc Tag with the EC50 of 3.1ng/ml determined by ELISA (QC Test).