

Human DLL4 Protein

Cat. No. DLL-HM104



Description

Source	Recombinant Human DLL4 Protein is expressed from HEK293 with His tag at the C-terminus. It contains Ser27-Pro524.
Accession	KT340-D
Molecular Weight	The protein has a predicted MW of 55.38 kDa. Due to glycosylation, the protein migrates to 56-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS, 200 mM L-Arginine (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Delta-like protein 4 (DLL4) is a type I membrane protein belonging to the Delta/Serrate/Lag2 (DSL) family of Notch ligands. Activates NOTCH1 and NOTCH4. Involved in angiogenesis; negatively regulates endothelial cell proliferation and migration and angiogenic sprouting. Essential for retinal progenitor proliferation. Required for suppressing rod fates in late retinal progenitors as well as for proper generation of other retinal cell types (By similarity). During spinal cord neurogenesis, inhibits V2a interneuron fate.

Assay Data


Bis-Tris PAGE



Human DLL4 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human DLL4 is greater than 90% as determined by SEC-HPLC.

ELISA Data

Human DLL4, His Tag ELISA

0.1µg Human DLL4, His Tag Per Well



Immobilized Human DLL4, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-DLL4 Antibody, hFc Tag with the EC50 of 1.6ng/ml determined by ELISA.