# Human DPPIV/CD26 Protein


## Cat. No. DPV-HM126

# ϗͶϲϿ·ႮႽ

Description	
Source	Recombinant Human DPPIV/CD26 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Asn29-Pro766.
Accession	P27487
Molecular Weight	The protein has a predicted MW of 86.39 kDa. Due to glycosylation, the protein migrates to 90-115 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	CD26/dipeptidyl peptidase (DPP)IV is a membrane-bound protein found in many cell types of the body, and a soluble form is present in body fluids. There is longstanding evidence that various primary tumors and also metastases express DPPIV/CD26 to a variable extent.


## Assay Data

#### **Bis-Tris PAGE**



Human DPPIV on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

## SEC-HPLC



The purity of Human DPPIV is greater than 95% as determined by SEC-HPLC.


# Human DPPIV/CD26 Protein


Cat. No. DPV-HM126

#### **Assay Data**

## **ELISA Data**

ϗͶϲϿ·ϽϨ





Immobilized Human DPPIV, His Tag at  $1\mu$ g/ml (100 $\mu$ I/well) on the plate. Dose response curve for Anti-DPPIV Antibody, hFc Tag with the EC50 of 4.2ng/ml determined by ELISA (QC Test)..

#### **Bioactivity Data**

Measured by its ability to cleave the fluorogenic peptide substrate, Gly-Pro-7-amido-4-methylcoumarin (GP-AMC). The specific activity is >8500 pmol/min/ $\mu$ g (QC Test).