Mouse ECSCR Protein

Cat. No. ECS-MM201



Description	
Source	Recombinant Mouse ECSCR Protein is expressed from HEK293 with hFc tag at the C-Terminus.
	It contains Gln19-Thr130.
Accession	Q3TZW0-1
Molecular Weight	The protein has a predicted MW of 38.6 kDa. Due to glycosylation, the protein migrates to 48-68 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	Storage

Formulation	Lyophilized from 0.22µm filtered solution in 20mM Tris,150mM NaCl (pH 8.2). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please


minimize freeze-thaw cycles.

Background

Endothelial cell-specific chemotaxis receptor (ECSCR) is a cell surface protein expressed by blood endothelial cells with roles in endothelial cell migration and signal transduction. Zebrafish ecscr is expressed in angioblasts and in axial vessels during angioblast migration and vasculogenesis. Morpholino-directed ecscr knockdown resulted in defective angioblast migration in the posterior lateral plate mesoderm, a process known to depend on vascular endothelial-derived growth factor (VEGF).

Assay Data

Bis-Tris PAGE




Mouse ECSCR on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. ECS-MM201



Assay Data



The purity of Mouse ECSCR is greater than 95% as determined by SEC-HPLC.