

# Cynomolgus EPHA2 Protein

Cat. No. EPH-CM4A2

## Description

<b>Source</b>	Recombinant Cynomolgus EPHA2 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Ala24-Ser534.
<b>Accession</b>	Q1KL86
<b>Molecular Weight</b>	The protein has a predicted MW of 59.2 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Ephrin type-A receptor 2 (EPHA2) is a receptor tyrosine kinase (RTK), whose over-expression has been observed in a variety of cancers, including breast cancer. EPHA2 expression may be causally related to tumorigenesis; therefore, it is important to understand how EPHA2 gene (EPHA2) expression is regulated.

## Assay Data


### Bis-Tris PAGE



Cynomolgus EPHA2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC

Assay Data



The purity of Cynomolgus EPHA2 is greater than 95% as determined by SEC-HPLC.