## Human EPHA4 Protein

## Cot No

## **KVCJ**US

Cat. No. EPH-HM1	
Description	
Source	Recombinant Human EPHA4 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Val20-Thr547.
Accession	P54764-1
Molecular Weight	The protein has a predicted MW of 59.5 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
Fully	> 95% as determined by HPLC
Formulation and Sto	rage
Formulation	Supplied as 0.22µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The expression and activation of EphA4 in the various cell types in a knee joint was upregulated upon an intraarticular injury. To determine if EphA4 signaling plays a role in osteoarthritis, we determined whether deficient EphA4 expression (in EphA4 knockout mice) or upregulation of the EphA4 signaling (with the EfnA4-fc treatment) would alter cellular functions of synoviocytes and articular chondrocytes. In synoviocytes, deficient EphA4 expression enhanced, whereas activation of the EphA4 signaling reduced, expression and secretion of key inflammatory cytokines and matrix metalloproteases.
Assay Data	
Bis-Tris PAGE MK R 140KD 115KD 80KD 70KD 50KD 40KD 30KD 25KD 15KD 10KD SEC-HPLC	Human EPHA4 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.
"AU 40 - 30 - 10 - 0 - 2 4	The purity of Human EPHA4 is greater than 95% as determined by SEC-HPLC.