

Mouse EPHA3 Protein

Cat. No. EPH-MM1A3

Description

| | |
|-------------------------|--|
| Source | Recombinant Mouse EPHA3 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Glu21-His541. |
| Accession | NP_034270.1 |
| Molecular Weight | The protein has a predicted MW of 59.7 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result. |
| Endotoxin | Less than 1EU per ug by the LAL method. |
| Purity | > 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC |

Formulation and Storage


| | |
|-----------------------|---|
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Erythropoietin-producing hepatocellular carcinoma cell surface type A receptor 3 (EPHA3) has been found to promote the proliferation and survival of prostate cancer (PCa) cell lines and prostate tumor development in nude mice. The interaction of AR and SP1 contributes to regulate EPHA3 expression, and the SP1 binding sites (295~261) in the EPHA3 core promoter region is crucial to the regulation of EPHA3 expression in response to androgen hormone stimuli.

Assay Data


Tris-Bis PAGE



Mouse EPHA3 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Mouse EPHA3 is greater than 95% as determined by SEC-HPLC.