

Biotinylated Human EPO R/Erythropoietin R Protein

Cat. No. EPO-HM50RB

Description

Source	Recombinant Biotinylated Human EPO R/Erythropoietin R Protein is expressed from HEK293 with hFc tag and Avi tag at the C-terminus. It contains Ala25-Pro250.
Accession	P19235-1
Molecular Weight	The protein has a predicted MW of 52.58 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The erythropoietin (epo) receptor is a member of the cytokine receptor family. It is expressed almost exclusively on erythroid precursor cells and controls the development of red blood cells. The epo receptor has no intrinsic kinase activity, but binds intracellular tyrosine kinases to elicit its signals. Alterations in the transmission of the signalling cascade lead to clinically abnormal red blood cell production.


Assay Data

Tris-Bis PAGE



Biotinylated Human EPO R on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Human EPO R is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Human EPOR, hFc Tag ELISA

0.5µg Human EPO, No Tag Per Well



Immobilized Human EPO, No Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human EPO R, hFc Tag with the EC50 of 0.30µg/ml determined by ELISA.