

# Cynomolgus FAP Protein

Cat. No. FAP-CM101



## Description

<b>Source</b>	Recombinant Cynomolgus FAP Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Leu26-Asp760.
<b>Accession</b>	XP_005573377
<b>Molecular Weight</b>	The protein has a predicted MW of 86.2 kDa. Due to glycosylation, the protein migrates to 80-100 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1EU per $\mu\text{g}$ by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

## Formulation and Storage


<b>Formulation</b>	Supplied as 0.22 $\mu\text{m}$ filtered solution in PBS, 20% Glycerol (pH 7.4).
<b>Storage</b>	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Fibroblast activation protein (FAP) is a serine protease that has been reported in fibroblasts and some carcinoma cells, which correlates with poor patient outcomes. FAP can be induced under hypoxia which is also vital in the malignant behaviors of cancer cells.


## Assay Data

### Bis-Tris PAGE



Cynomolgus FAP on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC




The purity of Cynomolgus FAP is greater than 95% as determined by SEC-HPLC.

## Assay Data

## ELISA Data

**Cynomolgus FAP, His Tag ELISA**

0.5µg Cynomolgus FAP, His Tag Per Well



Immobilized Cynomolgus FAP at 5µg/ml (100µl/well) on the plate. Dose response curve for Anti-FAP Antibody, hFc Tag with the EC50 of 5.3ng/ml determined by ELISA (QC Test).

## Bioactivity Data

Measured by its ability to convert the substrate benzyloxycarbonyl-Gly-Pro-7-amido-4-methylcoumarin (Z-GP-AMC) to Z-Gly-Pro and 7-amino-4-methylcoumarin (AMC). The specific activity is >2500 pmol/min/µg (QC Test).