

Mouse FAP Protein

Cat. No. FAP-MM101

Description

| | |
|-------------------------|---|
| Source | Recombinant Mouse FAP Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Leu26-Asp761. |
| Accession | P97321-1 |
| Molecular Weight | The protein has a predicted MW of 86.4 kDa. Due to glycosylation, the protein migrates to 90-100 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |

Formulation and Storage

| | |
|--------------------|--|
| Formulation | Supplied as 0.22 μm filtered solution in PBS, 20% Glycerol (pH 7.4). |
| Storage | Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Fibroblast activation protein (FAP) is a serine protease that has been reported in fibroblasts and some carcinoma cells, which correlates with poor patient outcomes. FAP can be induced under hypoxia which is also vital in the malignant behaviors of cancer cells.

Assay Data

Bis-Tris PAGE




Mouse FAP on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

ELISA Data

Assay Data

Mouse FAP, His Tag ELISA

0.2µg Mouse FAP, His Tag Per Well



Immobilized Mouse FAP, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Anti-FAP Antibody, hFc Tag with the EC50 of 6.0ng/ml determined by ELISA (QC Test).

Bioactivity Data

Measured by its ability to convert the substrate benzyloxycarbonyl-Gly-Pro-7-amido-4-methylcoumarin (Z-GP-AMC) to Z-Gly-Pro and 7-amino-4-methylcoumarin (AMC). The specific activity is >2000 pmol/min/µg (QC Test).