

# Mouse FAP Protein

Cat. No. FAP-MM101

## Description

<b>Source</b>	Recombinant Mouse FAP Protein is expressed from HEK293 with His tag at the N-Terminus. It contains Leu26-Asp761.
<b>Accession</b>	P97321-1
<b>Molecular Weight</b>	The protein has a predicted MW of 86.4 kDa. Due to glycosylation, the protein migrates to 90-100 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1EU per $\mu\text{g}$ by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22 $\mu\text{m}$ filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Fibroblast activation protein (FAP) is a serine protease that has been reported in fibroblasts and some carcinoma cells, which correlates with poor patient outcomes. FAP can be induced under hypoxia which is also vital in the malignant behaviors of cancer cells.

## Assay Data

### Bis-Tris PAGE




Mouse FAP on Bis-Tris PAGE under reduced conditions. The purity is greater than 95%.

### ELISA Data

#### Mouse FAP, His Tag ELISA

0.2 $\mu\text{g}$  Mouse FAP, His Tag Per Well



Immobilized Mouse FAP, His Tag at 2 $\mu\text{g}/\text{ml}$  (100 $\mu\text{l}/\text{well}$ ) on the plate. Dose response curve for Anti-FAP Antibody, hFc Tag with the EC50 of 6.0ng/ml determined by ELISA (QC Test).

## Mouse FAP Protein

Cat. No. FAP-MM101



### Assay Data

#### Bioactivity Data

Measured by its ability to convert the substrate benzyloxycarbonyl-Gly-Pro-7-amido-4-methylcoumarin (Z-GP-AMC) to Z-Gly-Pro and 7-amino-4-methylcoumarin (AMC). The specific activity is >2000 pmol/min/μg (QC Test).