

# Human FGFR2 beta (IIIb) Domain Protein

Cat. No. FGF-HM0BD

## Description

<b>Source</b>	Recombinant Human FGFR2 beta (IIIb) Domain Protein is expressed from HEK293 with Avi tag at the C-Terminus. It contains Pro253-Glu378.
<b>Accession</b>	P21802-3
<b>Molecular Weight</b>	The protein has a predicted MW of 15.49 kDa. Due to glycosylation, the protein migrates to 35-50 kDa based on Tris-Bis PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

## Formulation and Storage

<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Four distinct genes encoding closely related FGF receptors, FGF R1 - 4, are known. All four genes for FGF Rs encode proteins with an N-terminal signal peptide, three immunoglobulin (Ig)-like domains, an acid-box region containing a run of acidic residues between the IgI and IgII domains, a transmembrane domain and the split tyrosine-kinase domain. Multiple forms of FGF R1 - 3 are generated by alternative splicing of the mRNAs. A frequent splicing event involving FGF R1 and 2 results in receptors containing all three Ig domains, referred to as the alpha isoform, or only IgII and IgIII, referred to as the beta isoform.

## Assay Data

### Tris-Bis PAGE




Human FGFR2 beta (IIIb) Domain on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC

# Human FGFR2 beta (IIIb) Domain Protein

Cat. No. FGF-HM0BD

## Assay Data




The purity of Human FGFR2 beta (IIIb) Domain is greater than 95% as determined by SEC-HPLC.

## ELISA Data

### Human FGFR2 beta (IIIb) Domain, No Tag ELISA

0.05 $\mu$ g Human FGFR2 beta (IIIb) Domain, No Tag Per Well



Immobilized Human FGFR2 beta (IIIb) Domain, Avi Tag at 0.5 $\mu$ g/ml (100 $\mu$ l/Well) on the plate. Dose response curve for Anti-FGFR2 (IIIb) Antibody, hFc Tag with the EC50 of 1.6ng/ml determined by ELISA.