Biotinylated Human FGFR4 beta Protein

Cat. No. FGF-HM4RBB



Recombinant Biotinylated Human FGFR4 beta Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
It contains Pro152-Asp369.
P22455-1
The protein has a predicted MW of 26.90 kDa. Due to glycosylation, the protein migrates to 40-60 kDa based on Tris-Bis PAGE result.
Less than 1EU per μg by the LAL method.
> 95% as determined by Tris-Bis PAGE
> 90% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for


optimal storage. Please minimize freeze-thaw cycles.

Background

Fibroblast growth factor receptor 4 (FGF R4), also known as CD334, is a 110 kDa glycosylated transmembrane receptor tyrosine kinase. Tyrosine-protein kinase that acts as cell-surface receptor for fibroblast growth factors and plays a role in the regulation of cell proliferation, differentiation and migration, and in regulation of lipid metabolism, bile acid biosynthesis, glucose uptake, vitamin D metabolism and phosphate homeostasis. Required for normal down-regulation of the expression of CYP7A1, the rate-limiting enzyme in bile acid synthesis, in response to FGF19.


Assay Data

Tris-Bis PAGE



Biotinylated Human FGFR4 beta on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Human FGFR4 beta is greater than 90% as determined by SEC-HPLC.