

Biotinylated Human FGL1 Protein (Primary Amine Labeling)

Cat. No. FGL-HM211B

Description

Source	Recombinant Biotinylated Human FGL1 Protein (Primary Amine Labeling) is expressed from HEK293 with hFc tag at the N-Terminus. It contains Asp64-Asn305.
Accession	Q08830
Molecular Weight	The protein has a predicted MW of 54.8 kDa. Due to glycosylation, the protein migrates to 60-66 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 90% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 μm filtered solution in 20mM PB, 250mM NaCl (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Fibrinogen-like protein 1 (FGL-1) is a protein that is structurally related to fibrinogen. In humans, FLG-1 is encoded by the FGL1 gene. Fibrinogen-like protein 1 is a member of the fibrinogen family of proteins, which also includes fibrinogen, fibrinogen-like protein 2, and clotting factors V, VIII, and XIII. Fibrinogen-like Protein 1 is a major immune inhibitory ligand of LAG-3.


Assay Data

Tris-Bis PAGE



Biotinylated Human FGL1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Human FGL1 is greater than 90% as determined by SEC-HPLC.

ELISA Data

Assay Data

Biotinylated Human FGL1, hFc Tag ELISA

0.1 μ g Human LAG3, His Tag Per Well



Immobilized Human LAG3, His Tag at 1 μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Human FGL1, hFc Tag with the EC50 of 4.2ng/ml determined by ELISA.