

Biotinylated Cynomolgus/Rhesus macaque FcRn Protein

Cat. No. FRN-CM401B

Description

Source	Recombinant Biotinylated Cynomolgus/Rhesus macaque FcRn Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Ala24-Ser297(FCGRT) and Ile21-Met119(B2M).
Accession	Q8SPV9&Q8SPW0
Molecular Weight	The protein has a predicted MW of 46.5 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The neonatal Fc receptor (FCRN) is an approximately 45 kDa transmembrane glycoprotein with structural homology to MHC class I proteins. It is widely expressed in endothelial and epithelial cells and plays an important role in IgG homeostasis and antigen presentation by dendritic cells. FCGRT & B2M heterodimer protein (FcRn complex) consist of two subunits: p51, and p14, and forms an MHC class I-like heterodimer.

Assay Data


Tris-Bis PAGE



Biotinylated Cynomolgus/Rhesus macaque FcRn on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC


Assay Data



The purity of Biotinylated Cynomolgus/Rhesus macaque FcRn is greater than 95% as determined by SEC-HPLC.

Assay Data

SPR Data



Biotinylated Cynomolgus/Rhesus macaque FcRn, His Tag captured on CM5 Chip via Anti-His Antibody can bind Rituximab, hFc Tag with an affinity constant of 0.48 μ M as determined in SPR assay (Biacore T200).