Biotinylated Mouse FcRn Protein

Cat. No. FRN-MM401B



Description	
Source	Recombinant Biotinylated Mouse FcRn Protein is expressed from HEK293 His tag and Avi tag at the C-Terminus.
	It contains Ser22-Ser297 (FCGRT) and Ile21-Met119 (B2M).
Accession	P01887&Q61559
Molecular Weight	The protein has a predicted MW of 47.1 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation and Storage	
Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The neonatal Fc receptor (FCRN) is an approximately 45 kDa transmembrane glycoprotein with structural homology to MHC class I proteins. It is widely expressed in endothelial and epithelial cells and plays an important role in IgG homeostasis and antigen presentation by dendritic cells.FCGRT & B2M heterodimer protein (FcRn complex) consist of two subunits: p51, and p14, and forms an MHC class I-like heterodimer.


Assay Data

Tris-Bis PAGE



Biotinylated Mouse FcRn on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Biotinylated Mouse FcRn is greater than 95% as determined by SEC-HPLC.


Biotinylated Mouse FcRn Protein

Cat. No. FRN-MM401B



Assay Data

SPR Data



Biotinylated Mouse FcRn, His Tag captured on CM5 Chip via anti-his antibody can bind Trastuzumab with an affinity constant of 1.22 nM as determined in SPR assay (Biacore T200).