
### **Biotinylated Rat FcRn Protein**

### FRN-RM401B Cat. No.

Description	
Source	Recombinant Biotinylated Rat FcRn Protein is expressed from HEK293 with His tag and Avi Tag at the C- terminus.
	It contains Ala23-Ser298(FCGRT) and Ile21-Met119(B2M).
Accession	P13599(FCGRT)&P07151(B2M)
Molecular Weight	The protein has a predicted MW of 33.9 kDa (FCGRT) and 11.63 kDa (B2M). Due to glycosylation, the protein migrates to 45-60 kDa (FCGRT) and 12-14 kDa (B2M) based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	The neonatal Fc receptor (FCRN) is an approximately 45 kDa transmembrane glycoprotein with structural homology to MHC class I proteins. It is widely expressed in endothelial and epithelial cells and plays an important role in IgG homeostasis and antigen presentation by dendritic cells.FCGRT & B2M heterodimer protein (FcRn

role in IgG homeostasis and antigen presentation by dendritic cells.FCGRT & B2M heterodimer protein (FcRn complex) consist of two subunits: p51, and p14, and forms an MHC class I-like heterodimer.


# Assay Data



Biotinylated Rat FcRn on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

KAGTUS

## SEC-HPLC



The purity of Biotinylated Rat FcRn is greater than 95% as determined by SEC-HPLC.