Human FSHB Protein


Cat. No. FSH-HM201

_

ϗͶϲϿ·ႮႽ

Source Recombinant Human FSHB Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Asn19-Glu129. Accession P01225 P01225 Molecular Weight The protein has a predicted MW of 39.2 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result. Endotoxin Less than 1EU per μg by the LAL method. > 95% as determined by Tris-Bis PAGE
It contains Asn19-Glu129. Accession P01225 Molecular The protein has a predicted MW of 39.2 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result. Endotoxin Less than 1EU per μg by the LAL method. > 95% as determined by Tris-Bis PAGE
Molecular Weight The protein has a predicted MW of 39.2 kDa. Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result. Endotoxin Less than 1EU per μg by the LAL method. > 95% as determined by Tris-Bis PAGE
Weight Tris-Bis PAGE result. Endotoxin Less than 1EU per μg by the LAL method. > 95% as determined by Tris-Bis PAGE
> 95% as determined by Tris-Bis PAGE
> 95% as determined by Tris-Bis PAGE
Purity > 95% as determined by HPLC
Formulation and Storage
Formulation Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background
Fertility is dependent on follicle-stimulating hormone (FSH), a product of gonadotrope cells of the anterior pituitary gland. Hypothalamic gonadotropin-releasing hormone (GnRH) and intra-pituitary activins are regarded as the primary drivers of FSH synthesis and secretion. Both stimulate expression of the FSH beta subunit gene (Fshb), although the underlying mechanisms of GnRH action are poorly described relative to those of the activins.
Assay Data





Human FSHB on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

