

Biotinylated Human GCGR/Glucagon receptor Protein

Cat. No. GCR-HM40RB



Description

Source	Recombinant Biotinylated Human GCGR/Glucagon receptor Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Ala26-Lys136.
Accession	40RB
Molecular Weight	The protein has a predicted MW of 15.96 kDa. Due to glycosylation, the protein migrates to 30-50 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The pancreatic hormone glucagon activates the glucagon receptor (GCGR), a class B seven-transmembrane G protein-coupled receptor that couples to the stimulatory heterotrimeric G protein and provokes PKA-dependent signaling cascades vital to hepatic glucose metabolism and islet insulin secretion. The GCGR expressed at the plasma membrane is constitutively ubiquitinated and upon agonist-activation, internalized GCGRs are deubiquitinated at early endosomes and recycled via Rab4-containing vesicles.


Assay Data

Bis-Tris PAGE



Biotinylated Human GCGR on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Human GCGR is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Human GCGR, His Avi Tag ELISA

0.05µg Biotinylated Human GCGR, His Avi Tag Per Well



Immobilized Biotinylated Human GCGR, His Avi Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-GCGR Antibody, hFc Tag with the EC50 of 11.5ng/ml determined by ELISA.