

Biotinylated Cynomolgus Latent GDF-8 Protein (pro&latent)

Cat. No. GDF-CM408B

Description

Source	Recombinant Biotinylated Cynomolgus Latent GDF-8 Protein (pro&latent) is expressed from HEK293 with His tag and Avi tag at the N-terminus. It contains Gly19-Ser375.
Accession	Q95J86
Molecular Weight	The protein has a predicted MW of 43.45 kDa. Due to glycosylation, the protein migrates to 13-15 kDa, 38-41 kDa and 48-52 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Growth/differentiation factor 8 (GDF8), or myostatin, negatively regulates muscle mass. GDF8 is held in a latent state through interactions with its N-terminal prodomain. GDF8, like numerous TGF-β family members, is a disulfidelinked dimer that is synthesized as a precursor protein which requires cleavage by a furin-like protease to yield an N-terminal prodomain and a C-terminal mature, signaling domain.


Assay Data

Bis-Tris PAGE



Biotinylated Cynomolgus Latent GDF-8 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Cynomolgus Latent GDF-8 is greater than 90% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Cynomolgus Latent GDF-8, His Avi Tag ELISA

0.05µg Biotinylated Cynomolgus Latent GDF-8, His Avi Tag Per Well



Immobilized Biotinylated Cynomolgus Latent GDF-8, His Avi Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Apitegromab, hFc Tag with the EC50 of 2.8ng/ml determined by ELISA.