Biotinylated Human GPC3/Glypican 3 Protein





| Description | |
|---------------------|--|
| Source | Recombinant Biotinylated Human GPC3/Glypican 3 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. |
| | It contains Gln25-His559. |
| Accession | P51654-1 |
| Molecular Weight | The protein has a predicted MW of 63.6 kDa. Due to glycosylation, the protein migrates to 42 kDa, 68-80 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |
| | > 95% as determined by HPLC |
| | |

Formulation and Storage


| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before |
|----------------|---|
| | lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Glypican-3 is a protein ,which is encoded by the GPC3 gene in humans. The protein core of GPC3 consists of two subunits, where the N-terminal subunit has a size of ~40 kDa and the C-terminal subunit is ~30 kDa. Glypican 3 is a potential therapeutic target for treating liver cancer and other cancers. Several therapeutic anti-GPC3 antibodies have been developed.


Assay Data

Bis-Tris PAGE



Biotinylated Human GPC3 on Bis-Tris PAGE under reduced condition. The purity is greater than 90%.

SEC-HPLC




The purity of Biotinylated Human GPC3 is greater than 95% as determined by SEC-HPLC.

KAGTUS

Assay Data

ELISA Data

Biotinylated Human GPC3, His Tag ELISA 0.1μg Biotinylated Human GPC3, His Tag Per Well



Immobilized Biotinylated Human GPC3, His Tag at $1\mu g/ml$ ($100\mu l/well$) on the streptavidin precoated plate ($5\mu g/ml$). Dose response curve for Anti-GPC3 Antibody, hFc Tag with the EC50 of 6.4ng/ml determined by ELISA (QC Test).