
Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer Protein

| Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer Protein | | ϗͶϲͽ·υϩ |
|---|---|--|
| Cat. No. HLG-HM | 141C | |
| Description | | |
| Source | Recombinant Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. | |
| | It contains Gly25-Thr305(HLA-G), Ile21-Met119(B2M) and RIIPRHLQL peptide. | |
| Accession | P17693-1(HLA-G)&P61769(B2M)&RIIPRHLQL | |
| Molecular Weight | The protein has a predicted MW of 50.5 kDa. Due to glycosylation, the protein migrates to 51-60 kDa based on Bis-Tris PAGE result. | |
| Endotoxin | Less than 1 EU per μ g by the LAL method. | |
| Purity | > 95% as determined by Bis-Tris PAGE | |
| | > 95% as determined by HPLC | |
| Formulation and S | torage | |
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. | |
| Reconstitution | Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions. | |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 to aliquot the protein into smaller quantities for optimal storage. Please | |
| Background | | |
| | HLA-G is a molecule that was first known to confer protection to the fetu of its mother, thus critically contributing to fetal-maternal tolerance. The basis for HLA-G research and can be summarized as such: HLA-G, me its inhibitory receptors on immune cells (NK, T, B, monocytes/dendritic effectors, and so induces immune inhibition. | first functional finding constituted the mbrane-bound or soluble, strongly binds |

Assay Data

Bis-Tris PAGE




Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer Protein

Cat. No. HLG-HM41C

Assay Data

ĸΛϢ




The purity of Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer is greater than 95% as determined by SEC-HPLC.

ELISA Data

For Research Use Only





Immobilized Human HLA-G&B2M&Peptide (RIIPRHLQL) Monomer, His Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Anti-HLA-G Antibody, hFc Tag with the EC50 of 5.8ng/ml determined by ELISA.