

# Cynomolgus ICAM-1/CD54 Protein, Ultra Low Endotoxin

Cat. No. ICM-CM101-UL

## Description

<b>Source</b>	Recombinant Cynomolgus ICAM-1/CD54 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln28-Val480.
<b>Accession</b>	A0A2K5UKF7
<b>Molecular Weight</b>	The protein has a predicted MW of 50.55 kDa. Due to glycosylation, the protein migrates to 75-100 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 0.01 EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

## Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Intercellular adhesion molecule-1 (ICAM-1; CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis.

## Assay Data


### Bis-Tris PAGE



Cynomolgus ICAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

### SEC-HPLC

Assay Data



The purity of Cynomolgus ICAM-1 is greater than 95% as determined by SEC-HPLC.