

Cynomolgus ICAM-1/CD54 Protein

Cat. No. ICM-CM101

Description

Source	Recombinant Cynomolgus ICAM-1/CD54 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln28-Val480.
Accession	A0A2K5UKF7
Molecular Weight	The protein has a predicted MW of 50.55 kDa. Due to glycosylation, the protein migrates to 75-100 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Intercellular adhesion molecule-1 (ICAM-1; CD54) is a 90 kDa member of the immunoglobulin (Ig) superfamily and is critical for the firm arrest and transmigration of leukocytes out of blood vessels and into tissues. ICAM-1 is constitutively present on endothelial cells, but its expression is increased by proinflammatory cytokines. The endothelial expression of ICAM-1 is increased in atherosclerotic and transplant-associated atherosclerotic tissue and in animal models of atherosclerosis.

Assay Data


Tris-Bis PAGE



Cynomolgus ICAM-1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data



The purity of Cynomolgus ICAM-1 is greater than 95% as determined by SEC-HPLC.