

Mouse IFN alpha/beta R1 Protein

Cat. No. IFN-MM4R1

Description

Source	Recombinant Mouse IFN alpha/beta R1 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Glu27-Thr429.
Accession	P33896
Molecular Weight	The protein has a predicted MW of 46.9 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IFN-alpha / beta R1, also known as IFNAR1, belongs to the class II cytokine receptor family of proteins. Class II cytokine receptors form heterodimeric receptor complexes that mediate class II cytokine signals. Subunits of the different receptor complexes are shared and serve multiple functions. Functions in general as heterodimer with IFNAR2. Type I interferon binding activates the JAK-STAT signaling cascade, and triggers tyrosine phosphorylation of a number of proteins including JAKs, TYK2, STAT proteins and the IFNR alpha- and beta-subunits themselves.

Assay Data


Bis-Tris PAGE



Mouse IFNAR1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Assay Data



The purity of Mouse IFNAR1 is greater than 90% as determined by SEC-HPLC.

SPR Data



Mouse IFNAR1, His Tag captured on CM5 Chip via anti-his antibody can bind Human IFN alpha 1, hFc Tag with an affinity constant of 3.61 nM as determined in SPR assay (Biacore T200).