

Biotinylated Cynomolgus IgE Protein

Cat. No. IGE-CM401B

Description

Source	Recombinant Biotinylated Cynomolgus IgE Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Cys209-Lys429.
Accession	G8F4W7
Molecular Weight	The protein has a predicted MW of 27.9 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Immunoglobulin E (IgE) is well known for its role in allergic disease, the manifestations of which are mediated through its two Fc receptors, FcεRI and CD23 (FcεRII). IgE and its interactions with these receptors are therefore potential targets for therapeutic intervention, and exciting progress has been made in this direction. Furthermore, recent structural studies of IgE-Fc, the two receptors, and of their complexes, have revealed a remarkable degree of plasticity at the IgE-CD23 interface and an even more remarkable degree of dynamic flexibility within the IgE molecule.

Assay Data

Bis-Tris PAGE




Biotinylated Cynomolgus IgE on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Biotinylated Cynomolgus IgE Protein

Cat. No. IGE-CM401B

Assay Data



The purity of Biotinylated Cynomolgus IgE is greater than 95% as determined by SEC-HPLC.

Biotinylated Cynomolgus IgE Protein


Cat. No. IGE-CM401B

Assay Data

ELISA Data

Biotinylated Cynomolgus IgE, His Avi Tag ELISA

0.1µg Human Fc epsilon RI alpha, hFc Tag Per Well




Log Biotinylated Cynomolgus IgE, His Avi Tag Conc.(µg/ml)

Immobilized Human Fc epsilon RI alpha, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus IgE, His Avi Tag with the EC50 of 10.1ng/ml determined by ELISA.

ELISA Data

Biotinylated Cynomolgus IgE, His Avi Tag ELISA

0.05µg Biotinylated Cynomolgus IgE, His Avi Tag Per Well




Log Omalizumab, hFc Tag Conc.(µg/ml)

Immobilized Biotinylated Cynomolgus IgE, His Avi Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Omalizumab, hFc Tag with the EC50 of 5.1ng/ml determined by ELISA.

ELISA Data

Biotinylated Cynomolgus IgE, His Avi Tag ELISA

0.1µg Cynomolgus Fc epsilon RI alpha, His Tag Per Well



Log Biotinylated Cynomolgus IgE, His Avi Tag Conc.(µg/ml)

Immobilized Cynomolgus Fc epsilon RI alpha, His Tag (Cat. FER-CM101) at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus IgE, His Avi Tag with the EC50 of 8.8ng/ml determined by ELISA.