

Biotinylated Rhesus macaque IGFBP-7 Protein

Cat. No. IGF-CM4P7B



Description

Source	Recombinant Biotinylated Rhesus macaque IGFBP-7 Protein is expressed from HEK293 with His tag and Avi tag at the C-terminus.
	It contains Ser27-Leu282.
Accession	XP_001083041.3
Molecular Weight	The protein has a predicted MW of 29.83 kDa. Due to glycosylation, the protein migrates to 35-45 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IGFBP-7, also known as Mac25/Angiomodulin (AGM), GFBP-rp1, tumor-derived adhesion factor (TAF) and prostacyclin-stimulating factor (PSF), is a secreted protein that contains three protein domain modules. Human IGFBP-rp1 cDNA encodes 282 amino acid (aa) residue precursor protein with a putative 26 aa signal peptide. IGFBP-7 binds IGF-I and IGF-II with a relatively low affinity. Stimulates prostacyclin (PGI₂) production. Stimulates cell adhesion.

Assay Data

Bis-Tris PAGE



Biotinylated Rhesus macaque IGFBP-7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Rhesus macaque IGFBP-7 Protein

Cat. No. IGF-CM4P7B



Assay Data



The purity of Biotinylated Rhesus macaque IGFBP-7 is greater than 95% as determined by SEC-HPLC.