

## Mouse IGFBP-7 Protein

Cat. No. IGF-MM1BP

### Description

Source	Recombinant Mouse IGFBP-7 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Ser26-Leu281.
Accession	Q61581
Molecular Weight	The protein has a predicted MW of 27.5 kDa. Due to glycosylation, the protein migrates to 35-38 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

### Formulation and Storage

Formulation	Lyophilized from 0.22µm filtered solution in PBS, 150mM NaCl (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### Background

IGFBP-7, also known as Mac25/Angiomodulin (AGM), GFBP-rp1, tumor-derived adhesion factor (TAF) and prostacyclin-stimulating factor (PSF), is a secreted protein that contains three protein domain modules. Human IGFBP-rp1 cDNA encodes 282 amino acid (aa) residue precursor protein with a putative 26 aa signal peptide. IGFBP-7 binds IGF-I and IGF-II with a relatively low affinity. Stimulates prostacyclin (PGI<sub>2</sub>) production. Stimulates cell adhesion.

### Assay Data

#### Bis-Tris PAGE



Mouse IGFBP-7 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


#### SEC-HPLC

Mouse IGFBP-7 Protein

Cat. No.   IGF-MM1BP



Assay Data



The purity of Mouse IGFBP-7 is greater than 95% as determined by SEC-HPLC.