

Human IL-13 Protein

Cat. No. IL1-HM213



Description

Source	Recombinant Human IL-13 Protein is expressed from HEK293 with hFc tag at the N-Terminus. It contains Gly21-Asn132.
Accession	AAK53823
Molecular Weight	The protein has a predicted MW of 39.6 kDa. Due to glycosylation, the protein migrates to 50-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin-13 (IL-13) is a monomeric 17 kDa immunoregulatory cytokine that plays a key role in the pathogenesis of allergy, cancer, and tissue fibrosis. It is secreted by several helper T cell subsets, NK cells, mast cells, eosinophils, basophils, and visceral smooth muscle cells. Inhibits inflammatory cytokine production. Synergizes with IL2 in regulating interferon-gamma synthesis. May be critical in regulating inflammatory and immune responses. Positively regulates IL31RA expression in macrophages (By similarity).

Assay Data


Bis-Tris PAGE



Human IL-13 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human IL-13 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human IL-13, hFc Tag, His Tag ELISA


0.2µg Biotinylated Human IL-13Ra1, His Tag Per Well



Immobilized Biotinylated Human IL-13Ra1, His Tag at 2µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human IL-13, hFc Tag with the EC50 of 15.8ng/ml determined by ELISA (QC Test).


ELISA Data

Assay Data

Human IL-13, hFc Tag ELISA0.1 μ g Human IL-13, hFc Tag Per Well

Immobilized Human IL-13, hFc Tag at 1 μ g/ml (100 μ l/well) on the plate. Dose response curve for Human IL-13Ra2, His Tag with the EC50 of 7.1ng/ml determined by ELISA.

SPR Data



Human IL-13, hFc Tag captured on CM5 Chip via Protein A can bind Human IL-13Ra1, His Tag with an affinity constant of 0.399 μ M as determined in SPR assay (Biacore T200).