

Biotinylated Human IL-10 Protein

Cat. No. IL1-HM410B

Description

Source	Recombinant Biotinylated Human IL-10 Protein is expressed from HEK293 with His tag and Avi tag at the N-Terminus. It contains Ser19-Asn178.
Accession	P22301
Molecular Weight	The protein has a predicted MW of 21.3 kDa. Due to glycosylation, the protein migrates to 23-25 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin 10 (IL10) is a key anti-inflammatory cytokine that can inhibit proinflammatory responses of both innate and adaptive immune cells. An association between IL10 and intestinal mucosal homeostasis became clear with the discovery that IL10 and IL10 receptor (IL10R)-deficient mice develop spontaneous intestinal inflammation. Similarly, patients with deleterious mutations in IL10, IL10RA, or IL10RB present with severe enterocolitis within the first months of life.


Assay Data

Bis-Tris PAGE



Biotinylated Human IL-10 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SPR Data



Human IL-10 R alpha, His Tag immobilized on CM5 Chip can bind Biotinylated Human IL-10, His Tag with an affinity constant of 0.60 nM as determined in SPR assay (Biacore T200).