Mouse IL-20RA Protein

Cat. No. IL2-MM2RA



| Description | |
|---------------------|--|
| Source | Recombinant Mouse IL-20RA Protein is expressed from HEK293 with hFc tag at the C-Terminus. |
| | It contains Val33-Lys253. |
| Accession | Q6PHB0 |
| Molecular Weight | The protein has a predicted MW of 52.1 kDa. Due to glycosylation, the protein migrates to 65-70 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE |
| | |

Formulation and Storage


| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
|----------------|---|
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

IL-20RA was highly expressed in the tumor tissue of CRC and related to the advanced stage.IL-20RA was involved in regulating oncogenic and immune pathways and affecting the expression of genes related to cell proliferation and immune evasion in CRC.Super-enhancer inhibition by JQ-1 or iBET-151 suppressed the growth of tumor cells and inhibited the expression of IL-20RA.


Assay Data

Bis-Tris PAGE



Mouse IL-20RA on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SPR Data



Mouse IL-20RA, hFc Tag captured on CM5 Chip via Protein A can bind Human IL-20, His Tag with an affinity constant of 13.93 μ M as determined in SPR assay (Biacore T200).