

Human IL-3 Protein

Cat. No. IL3-HE401



Description

Source	Recombinant Human IL-3 Protein is expressed from E.coli with His tag and Avi tag at the N-Terminus.
	It contains Ala20-Phe152.
Accession	P08700
Molecular Weight	The protein has a predicted MW of 16.9 kDa same as Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin 3 is a pleiotropic factor produced primarily by activated T cells that can stimulate the proliferation and differentiation of pluripotent hematopoietic stem cells as well as various lineage committed progenitors. This CSF induces granulocytes, macrophages, mast cells, stem cells, erythroid cells, eosinophils and megakaryocytes.

Assay Data

Bis-Tris PAGE



Human IL-3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Human IL-3 Protein

Cat. No. IL3-HE401




Assay Data



The purity of Human IL-3 is greater than 95% as determined by SEC-HPLC.


ELISA Data

Human IL-3, His Tag ELISA
0.5µg Human IL-3, His Tag Per Well



Immobilized Human IL-3, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Human IL-3 R alpha, hFc Tag with the EC50 of 0.46µg/ml determined by ELISA (QC Test).

Cell Based Assay



Measured in a cell proliferation assay using TF1 human erythroleukemic cells. The ED50 for this effect is < 0.2 ng/mL.