

Human IL-7 R alpha/CD127 Protein, Ultra Low Endotoxin



Cat. No. IL7-HM2RA-UL

Description	
Source	Recombinant Human IL-7 R alpha/CD127 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Glu21-Gly236.
Accession	P16871-1
Molecular Weight	The protein has a predicted MW of 51.6 kDa. Due to glycosylation, the protein migrates to 65-75 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.01 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Interleukin 7 (IL-7) and its receptor (IL-7R, a heterodimer of IL-7Rα and γc) are essential for normal lymphoid development. IL-7 and IL-7R activate three main pathways: STAT5, PI3K/Akt/mTOR and MEK/Erk, ultimately leading to the promotion of leukemia cell viability, cell cycle progression and growth.	

Assay Data


Bis-Tris PAGE



Human IL-7 R alpha on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human IL-7 R alpha is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human IL-7R alpha, hFc Tag ELISA

0.2µg Human IL-7, No Tag Per Well



Immobilized Human IL-7 at 2µg/ml (100µl/Well) on the plate. Dose response curve for Human IL-7 R alpha, hFc Tag with the EC50 of 16.2ng/ml determined by ELISA (QC tested).