

Cynomolgus IL-18 R1/CD218a Protein

Cat. No. IL8-CM1R1

Description

Source	Recombinant Cynomolgus IL-18 R1/CD218a Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Cys22-Gly329.
Accession	A0A2K5V3J9
Molecular Weight	The protein has a predicted MW of 36.45 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 μm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin-18 (IL-18) is a member of the interleukin-1 family of cytokines produced constitutively by different cell types and by adipose tissue. Interleukin (IL)-18 was originally discovered as a factor that enhanced IFN- γ production from anti-CD3-stimulated Th1 cells, especially in the presence of IL-12. Upon stimulation with Ag plus IL-12, naïve T cells develop into IL-18 receptor (IL-18R) expressing Th1 cells, which increase IFN- γ production in response to IL-18 stimulation.

Assay Data


Bis-Tris PAGE



Cynomolgus IL-18 R1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Assay Data



The purity of Cynomolgus IL-18 R1 is greater than 95% as determined by SEC-HPLC.

SPR Data



Cynomolgus IL-18 R1, His Tag captured on CM5 Chip via anti-his antibody can bind Human IL-18, No Tag with an affinity constant of 2.02 nM as determined in SPR assay (Biacore T200).