## Mouse IL-18RAP Protein

## ᠺ᠕ᡗ᠋ᠴᡃ᠐᠍᠋

## Cat. No. IL8-MM1AP Description Recombinant Mouse IL-18RAP Protein is expressed from HEK293 with His tag at the C-Terminus. Source It contains Phe20-Glu356. Accession Q9Z2B1 Molecular The protein has a predicted MW of 39.7 kDa. Due to glycosylation, the protein migrates to 55-65 kDa based on Weight Tris-Bis PAGE result. Endotoxin Less than 1EU per µg by the LAL method. > 95% as determined by Tris-Bis PAGE Purity > 95% as determined by HPLC Formulation and Storage Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before Formulation lyophilization. Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Reconstitution Dissolve the lyophilized protein in distilled water. -20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3-6 months after reconstitution.2-8°C for 2-7 Storage days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. Background lactoferrin (LTF), peptidoglycan recognition protein 1 and IL-18 receptor accessory protein (IL18RAP) remained significantly elevated in peripheral monocytes of hypertensive individuals in a separate validation cohort. Interestingly, IL18RAP expression associated with MAP in a cohort of African Americans.. Assay Data **Tris-Bis PAGE** MK R 140KD 115KD 80KD 70KD 50KD Mouse IL-18RAP on Tris-Bis PAGE under 40KD reduced condition. The purity is greater than 95%. 30KD 25KD 15KD 10KD SEC-HPLC m A U 50 40 The purity of Mouse IL-18RAP is greater than 30 95% as determined by SEC-HPLC. 20

10