

# Mouse IL-18 R1/CD218a Protein

Cat. No. IL8-MM1R1

## Description

<b>Source</b>	Recombinant Mouse IL-18 R1/CD218a Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Lys20-Gly326.
<b>Accession</b>	Q61098
<b>Molecular Weight</b>	The protein has a predicted MW of 36.5 kDa. Due to glycosylation, the protein migrates to 60-80 kDa based on Tris-Bis PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

## Formulation and Storage

<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Interleukin-18 (IL-18) is a member of the interleukin-1 family of cytokines produced constitutively by different cell types and by adipose tissue. Interleukin (IL)-18 was originally discovered as a factor that enhanced IFN-γ production from anti-CD3-stimulated Th1 cells, especially in the presence of IL-12. Upon stimulation with Ag plus IL-12, naive T cells develop into IL-18 receptor (IL-18R) expressing Th1 cells, which increase IFN-γ production in response to IL-18 stimulation.

## Assay Data


### Tris-Bis PAGE



Mouse IL-18 R1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


### SEC-HPLC

Assay Data



The purity of Mouse IL-18 R1 is greater than 95% as determined by SEC-HPLC.

SPR Data



Mouse IL-18 R1, His Tag captured on CM5 Chip via anti-his antibody can bind Human IL-18, No Tag with an affinity constant of 0.30  $\mu$ M as determined in SPR assay (Biacore T200).