

## Mouse IL-9 Protein

Cat. No. IL9-MM101

### Description

<b>Source</b>	Recombinant Mouse IL-9 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gln19-Pro144.
<b>Accession</b>	P15247
<b>Molecular Weight</b>	The protein has a predicted MW of 15.4 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Bis-Tris PAGE result.
<b>Endotoxin</b>	Less than 1EU per µg by the LAL method.
<b>Purity</b>	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

### Formulation and Storage


<b>Formulation</b>	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
<b>Reconstitution</b>	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
<b>Storage</b>	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 4-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

### Background

IL-9 is a pleiotropic cytokine that influences various distinct functions of different target cells such as T cells, B cells, mast cells and airway epithelial cells by activating STAT1, STAT3 and STAT5. Because of its pleiotropic functions, IL-9 has been demonstrated to be involved in several diseases, such as cancer, autoimmunity and other pathogen-mediated immune-regulated diseases. In this review, we focus on the role of Th9 and IL-9-producing cells in allergic asthma.

### Assay Data

#### Bis-Tris PAGE




Mouse IL-9 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

#### SEC-HPLC

# Mouse IL-9 Protein

Cat. No. IL9-MM101


## Assay Data



The purity of Mouse IL-9 is greater than 95% as determined by SEC-HPLC.

## Cell Based Assay

### Recombinant Mouse IL-9 Bioactivity



Measured in a cell proliferation assay using MO7e human megakaryocytic leukemic cells. The ED50 for this effect is < 3 ng/mL.