

# Mouse IL-17A/CTLA-8 Protein

Cat. No. ILA-MM417

KACUS

## Description

Source	Recombinant Mouse IL-17A/CTLA-8 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Ala26-Ala158.
Accession	Q62386-1
Molecular Weight	The protein has a predicted MW of 18 kDa. Due to glycosylation, the protein migrates to 20-28 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE

## Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

## Background

Interleukin17A (IL17A), also known as CTLA8, is a 1520 kDa glycosylated cytokine that plays an important role in antimicrobial and chronic inflammation. The six IL17 cytokines (IL17AF) are encoded by separate genes but adopt a conserved cystine knot fold. IL-17A is a ligand for IL17RA and IL17RC. The heterodimer formed by IL17A and IL17F is a ligand for the heterodimeric complex formed by IL17RA and IL17RC. Involved in inducing stromal cells to produce proinflammatory and hematopoietic cytokines.


## Assay Data

### Bis-Tris PAGE



### ELISA Data

**Mouse IL-17A, His Tag ELISA**  
0.2µg Mouse IL-17A, His Tag Per Well



Immobilized Mouse IL-17A, His Tag at 2µg/ml (100µl/well) on the plate. Dose response curve for Mouse IL-17R alpha, hFc Tag with the EC50 of 29.6ng/ml determined by ELISA (QC Test).