

Biotinylated Human IL-23 alpha&Mouse IL-12 beta Protein

Cat. No. ILB-HM4ABB

Description

Source	Recombinant Biotinylated Human IL-23 alpha&Mouse IL-12 beta Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Arg20-Pro189 (Human IL-23 alpha) and Met23-Ser335 (Mouse IL-12 beta).
Accession	Q9NPF7(Human IL-23 alpha)&P43432-1(Mouse IL-12 beta)
Molecular Weight	The protein has a predicted MW of 21.7 kDa (Human IL-23 alpha) & 35.8 kDa (Mouse IL-12 beta). Due to glycosylation, the protein migrates to 22-25 kDa (Human IL-23 alpha) & 45-60 kDa (Mouse IL-12 beta) based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin (IL)-12 and IL-23 belong to the IL-12 type family and are composite cytokines, consisting of the common β subunit p40 and the specific cytokine α subunit p35 and p19, respectively. IL-12 signals via the IL-12R β 1·IL-12R β 2 receptor complex, and IL-23 uses also IL-12R β 1 but engages IL-23R as second receptor.

Assay Data


Tris-Bis PAGE



Biotinylated Human IL-23 alpha&Mouse IL-12 beta on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Biotinylated Human IL-23 alpha&Mouse IL-12 beta is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Human IL-23 alpha & Mouse IL-12 beta, His Tag ELISA

0.1µg Biotinylated Human IL-23 alpha & Mouse IL-12 beta, His Tag Per Well



Immobilized Biotinylated Human IL-23 alpha & Mouse IL-12 beta, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human IL-23R, mFc Tag with the EC50 of 28.1ng/ml determined by ELISA.