

Human ILDR2 Protein

Cat. No. ILD-HM202

Description

| | |
|-------------------------|--|
| Source | Recombinant Human ILDR2 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Leu21-Met184. |
| Accession | Q71H61 |
| Molecular Weight | The protein has a predicted MW of 35 kDa. Due to glycosylation, the protein migrates to 48-55 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per µg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 90% as determined by HPLC |

Formulation and Storage

| | |
|--------------------|--|
| Formulation | Supplied as 0.22µm filtered solution in PBS (pH 7.4). |
| Storage | Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Ildr2, a modifier of diabetes susceptibility in obese mice, is expressed in most organs, including islets and hypothalamus, with reduced levels in livers of diabetes-susceptible B6.DBA mice congenic for a 1.8 Mb interval of Chromosome 1. In hepatoma and neuronal cells, ILDR2 is primarily located in the endoplasmic reticulum membrane. Livers in knockdown mice were steatotic, with increased hepatic and circulating triglycerides and total cholesterol. Increased circulating VLDL, without reduction in triglyceride clearance suggests an effect of reduced hepatic ILDR2 on hepatic cholesterol clearance.


Assay Data

Bis-Tris PAGE



Human ILDR2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human ILDR2 is greater than 90% as determined by SEC-HPLC.