

Canine IL-23R Protein

Cat. No. ILR-DM123

Description

Source	Recombinant Canine IL-23R Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Gly24-Asp354.
Accession	A0A8I3PXL0
Molecular Weight	The protein has a predicted MW of 39.31 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Supplied as 0.22 μm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Interleukin-23 (IL-23), a member of the IL-12 family of cytokines, is a heterodimeric cytokine. It is composed of subunits p40 (shared with IL-12) and p19 (an IL-12 p35-related subunit) and is secreted by several types of immune cells, such as natural killer cells and dendritic cells. The IL-23 receptor is composed of the subunit IL-12R β 1 and the IL-23-specific subunit IL-23R. The binding of IL-23 to its specific cell surface receptor regulates a number of functions, including proliferation and differentiation of cells and secretion of cell factors.


Assay Data

Bis-Tris PAGE



Canine IL-23R on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Canine IL-23R is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Canine IL-23R, His Tag ELISA

0.5µg Canine IL-23R, His Tag Per Well



Immobilized Canine IL-23R, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human IL-23 alpha&IL-12 beta, His Tag with the EC50 of 2.7µg/ml determined by ELISA.

Log Biotinylated Human IL-23 alpha&IL-12 beta, His Tag Conc.(µg/ml)