

Biotinylated Human IL-22R alpha 1&IL-10R beta Protein

Cat. No. ILR-HM5ABB

Description

Source	Recombinant Biotinylated Human IL-22R alpha 1&IL-10R beta Protein is expressed from HEK293 with hFc tag and Avi Tag at the C-Terminus. It contains His16-Thr228 (IL-22R alpha 1) and Met20-Ser220 (IL-10R beta).
Accession	Q8N6P7(IL-22R alpha 1)&Q08334(IL-10R beta)
Molecular Weight	The protein has a predicted MW of 52.07 kDa (IL-22R alpha 1) and 48.85 kDa (IL-10R beta). Due to glycosylation, the protein migrates to 68-75 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Supplied as 0.22µm filtered solution in PBS (pH 7.4).
Storage	Valid for 12 months from date of receipt when stored at -80°C. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

IL-22R α 1&IL-10R β are receptors for IL-22 and IL-10, mediate the activation of JAK-STAT, MAPK and other signaling pathways, and participate in the process of many inflammatory diseases. IL-22 R α 1 contains cytoplasmic motifs for interactions with signal transduction molecules, but formation of ternary complexes with IL10 R β or IL20 R β and the respective ligands is required for signal transduction[1].

Assay Data

Bis-Tris PAGE



Biotinylated Human IL-22R alpha 1&IL-10R beta on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human IL-22R alpha 1&IL-10R beta Protein

Cat. No. ILR-HM5ABB



Assay Data




The purity of Biotinylated Human IL-22R alpha 1&IL-10R beta is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human IL-22R alpha 1&IL-10R beta, hFc Tag ELISA

0.5µg Human IL-22, His Tag Per Well



Immobilized Human IL-22, His Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Human IL-22R alpha1&IL-10R beta, hFc Tag with the EC50 of 1.36µg/ml determined by ELISA.

Log Biotinylated Human IL-22R alpha 1&IL-10R beta, hFc Tag Conc. (µg/ml)